

24 July 1996

Mesospheric Response to Impacting Relativistic Electrons

Final Report

Period: 21 July 1995 through 20 July 1996

Contract P.O. S-57779-F

D.L. Chenette
Lead Investigator

Space Sciences Laboratory
Lockheed-Martin Palo Alto Research Laboratories
LMMS/ATC O/91-20 B/252
Palo Alto, CA 94304-1191

24 July 1996

Mesospheric Response to Impacting Relativistic Electrons

Final Report

Period: 21 July 1995 through 20 July 1996

Contract P.O. S-57779-F

D.L. Chenette
Lead Investigator

Space Sciences Laboratory
Lockheed-Martin Palo Alto Research Laboratories
LMMS/ATC O/91-20 B/252
Palo Alto, CA 94304-1191



Report Documentation Page

1. Report No.	2. Government Accession No.	3. Recipient's Catalog No.
4. Title and Subtitle Mesospheric Response to Impacting Relativistic Electrons-Final Report		5. Report Date 24 July 1996
7. Author(s) D. L. Chenette		6. Performing Organization Code
9. Performing Organization Name and Address LMMS/ATC, O/H1-11, B/252 3251 Hanover St. Palo Alto, CA 94304		8. Performing Organization Report No.
12. Sponsoring Agency Name and Address NASA/Goddard Space Flight Center Greenbelt, MD 20771		10. Work Unit No.
15. Supplementary Notes		11. Contract or Grant No. P.O. S-57779-F
		13. Type of Report and Period Covered 21 July 1995-Final Report 20 July 1996
16. Abstract Daily maps of the spatial distributions of precipitating energetic electrons were produced for the period 10-20 May 1992. These data will serve as the input for potential changes in upper atmosphere composition.		14. Sponsoring Agency Code
17. Key Words (Suggested by Author(s)) mesosphere; relativistic electrons; electron precipitation; ion production; UARS; PEM; HEPS		18. Distribution Statement
19. Security Classif. (of this report)	20. Security Classif. (of this page)	21. No. of pages
U	U	3+4 Apps
22. Price		

REPORT DOCUMENTATION PAGE

Form Approved:
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20501.

1. AGENCY USE ONLY (Leave blank)			2. REPORT DATE 24 July 1996		3. REPORT TYPE AND DATES COVERED Final Report 21 July '95-20 July '96		
4. TITLE AND SUBTITLE Mesospheric Response to Impacting Relativistic Electrons-Final Report			5. FUNDING NUMBERS P.O. S-57779-F				
6. AUTHOR(S) D. L. Chenette							
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) LMMS/ATC, O/H1-11, B/252 3251 Hanover St. Palo Alto, CA 94304			8. PERFORMING ORGANIZATION REPORT NUMBER				
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) NASA/Goddard Space Flight Center Greenbelt, MD 20771			10. SPONSORING/MONITORING AGENCY REPORT NUMBER				
11. SUPPLEMENTARY NOTES							
12a. DISTRIBUTION/AVAILABILITY STATEMENT				12b. DISTRIBUTION CODE			
13. ABSTRACT (Maximum 200 words) Daily maps of the spatial distributions of precipitating energetic electrons were produced for the period 10-20 May 1992. These data will serve as the input for potential changes in upper atmosphere composition.							
14. SUBJECT TERMS mesosphere; relativistic electrons; electron precipitation; ion production; UARS; PEM; HEPS					15. NUMBER OF PAGES 3+4 Apps		
					16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT U		18. SECURITY CLASSIFICATION OF THIS PAGE U		19. SECURITY CLASSIFICATION OF ABSTRACT U		20. LIMITATION OF ABSTRACT	

Introduction

The Space Sciences Laboratory (O/H1-11) of the Lockheed Martin Missiles and Space Company's Advanced Technology Center (LMMS/ATC) is supporting Dr. Richard Goldberg of the NASA Goddard Space Flight Center in a scientific investigation of the effects of relativistic electron precipitation on the chemical composition and dynamics of the mesosphere. This is the final progress report of the 1-year effort performed at LMMS in this collaboration.

Results

During the past year most of the resources provided by the purchase order were used to support the efforts of Mr. Edward E. Gaines and Spacerad in the reduction and analysis of the electron data from the PEM/HEPS instrument aboard UARS. As a result, the bulk of this report consists of Mr. Gaines's reports of his progress and results. These are attached as Appendix A, B, and C.

The results described in the Appendix have been provided informally to Dr. Goldberg and discussed with him in face-to-face meetings at Lockheed Martin. In addition, daily maps of precipitating electron intensity have been produced for the May 1992 period of highest precipitating electron intensity. These also have been provided directly to Dr. Goldberg. Copies are attached.

Summary

Based on plots of the daily electron intensity, the largest events observed aboard UARS for the period from October 1991 through mid-1995 were identified. Based on these results together with careful consideration of the availability of data from other UARS investigations, a prioritized list of intervals was agreed upon for more detailed investigation. The following intervals were selected, shown here in priority order:

11 - 21 May 1992
10 - 20 October 1991
24 November - 21 December 1991
4 - 8 November 1991

and 6 April - 3 May 1993


The 11-21 May 1992 interval is the highest priority and in discussions with Dr. Goldberg we agreed to give it the most attention in our analysis.


Relativistic electron energy spectra from the HEPS instrument were calculated in detail for each day of the May 1992 period, and in summary form for the November 1991 and December 1991 periods. Indications of the temporal and spatial variations of the precipitating electron intensity were provided with samples obtained at different places and times within each event period. The precipitated electron flux intensities and energy spectra were also used to calculate altitude profiles of ion production rates throughout the mesosphere. During the most intense parts of the May 1992 period, between the 12th and 14th of May, ion production rates exceeded 100 ions $\text{cm}^{-3} \text{ s}^{-1}$ at altitudes between 55 and 80 km.


Daily maps of the precipitating electron flux intensity were produced for the May 1992 period. They showed that the most intense precipitation observed by UARS was in the southern hemisphere, south of Africa and between 45° and 60° south magnetic latitude. This localization effect is at least partially the result of the geometry of the earth's magnetic field, specifically the offset of the magnetic center towards the north Pacific region.


Conclusions and Recommendations


Based on this study we conclude that there were extended periods of significant enhancements in the ion production rate down to below 60 km altitude as a result of relativistic electron precipitation. This result is quantified in the attached reports. The major electron precipitation event of May 1992 is thoroughly analyzed. We recommend that the spatial and temporal variations in the electron precipitation be compared to variations in the composition of trace species as revealed by the UARS atmospheric composition instruments, specifically CLAES and HALOE, but possibly also including MLS. We look forward to a continuation of this collaboration for two more years to work on this aspect of the problem. Now that we have characterized the input, in this way we hope to characterize the mesospheric response of the impacting relativistic electrons.
























































HEPS ELECTRONS 20 MAY 1992 HEPS 2 HEAD 1: TRAPPED ELECTRONS E>1 MEV




HEPS ELECTRONS 21 MAY 1992 HEPS 1 HEAD 2: PRECIPITATING ELECTRONS $E > 1$ MEV




HEPS ELECTRONS 21 MAY 1992 HEPS 2 HEAD 1: TRAPPED ELECTRONS $E > 1$ MEV



HEPS ELECTRONS 22 MAY 1992 HEPS 1 HEAD 2: PRECIPITATING ELECTRONS $E > 1$ MEV



HEPS ELECTRONS 22 MAY 1992 HEPS 2 HEAD 1: TRAPPED ELECTRONS $E > 1$ MEV



Appendix A

30 Nov. 1995

Interim Report - Mesospheric Response to Impacting Relativistic

Electrons : Task 1. Ref P.O. SC TDZ0930 F


Submitted in completion of Task 1 of SPACERAD statement of work:


Using data from the PEM/HEPS instrument aboard the UARS spacecraft, periods of highly relativistic electron precipitation will be identified, and SPACERAD will join in the selection of specific periods for more detailed analysis.

The intensity of relativistic electron fluxes at the UARS spacecraft has been extracted for the time period from instrument turn-on to mid-April 1995 when continuous operation of the HEPS was interrupted due to spacecraft difficulties. Plots of the peak daily flux of electrons with energies greater than 2 MeV are attached.

From the plots, several periods of high electron flux intensity were selected for suggested study during the operational lifetime of the CLAES instrument with which correlations of electron precipitation are to be sought. A summary of these suggested intervals is also attached.


— FLUX>2MEV






— FLUX>2MEV


HEPS ELECTRON FLUX - E > 2 MEV, DAILY PEAK VALUE - 1993





— FLUX > 2 MeV

HEPS ELECTRON FLUX - E > 2 MEV, DAILY PEAK VALUE - 1995



Mesospheric Response to Impacting Relativistic Electrons

SUGGESTED INTERVALS OF HRE FLUX FOR DETAILED STUDY

First priority - with CLAES data

DATE	UARS DAYS	PEAK > 2 MEV TRAP FLUX
1. May 11-21, 1992	243 - 253	2. X 10 ⁴ / cm ² -s-sr.
2. Oct. 10-20, 1991	29 - 39	8. X 10 ³
3. Nov. 24-Dec. 21, 1991	74 - 101	7. X 10 ³
4. Nov. 4-8, 1991	54 - 58	6. X 10 ³
5. Apr. 6-May 3, 1993	573 - 600	6. X 10 ³

Appendix B

Interim Report


26 Feb. 1996

Task 2 in Statement of Work for support of project:
Mesospheric Response to Impacting Relativistic Electrons
Ref. P.O.# SC PDZ 0931F


Energy spectra, spatial and temporal distributions of precipitating relativistic electrons and resulting ion production profiles have been obtained from PEM/HEPS data in the time periods identified in Task 1 of this effort as most promising for correlation with measurements of ozone by other instruments on board the UARS spacecraft. These results have been transmitted to LMMS Dept. 91-20 and to Dr. R. A. Goldberg of NASA, and figures illustrating the measurements are attached herewith.

The periods for detailed analysis include the most intense fluxes of relativistic electrons observed with the HEPS instrument (May 12 - 20, 1992) and other periods with large changes in the fluxes of precipitating relativistic electrons within a few days. Times of both northward and southward observation by CLAES are included, but the magnetic local times of the flux observation periods were all between 0900 and 2000 hours.


PRECIPITATING ELECTRON SPECTRA, 8-9 NOV. 1991, SOUTH




D 96 - 97 PRECIPITATING ELECTRONS : DLC REGION




ION PRODUCTION PROFILES, 8 - 9 NOV., 16 - 17 DEC. 1991, 8 & 12 MAY 1992




BACKGROUND ELECTRON SPECTRA, 9 NOV. 1991 & MAY 1992




PRECIPITATING ELECTRON SPECTRA, 12 MAY 1992




PRECIPITATING ELECTRON SPECTRA 13 MAY 1992, 23:14 - 23:26 UT




PRECIPITATING ELECTRON SPECTRA 14 MAY, 1992 - 21:40-21:52 UT




PRECIPITATING ELECTRON SPECTRA, 15 MAY 1992 - 18:29-18:31 UT




PRECIPITATING ELECTRON SPECTRA 16 MAY 1992, 20:19 - 20:23 UT




PRECIPITATING ELECTRON SPECTRA 17 MAY 1992, 18:41 - 18:49 UT




PRECIPITATING ELECTRON SPECTRA 18 MAY, 1992 - 23:31-23:38 UT



PRECIPITATING ELECTRON SPECTRA 19 MAY 1992 , 20:29 - 20:33 UT



ION PRODUCTION PROFILES FROM RELATIVISTIC ELECTRONS, MAY 1992



Appendix C

1996 Final Report

Mesospheric Response to Impacting Relativistic Electrons

Introduction

The purpose of this investigation is to search for correlation between precipitating high-energy electrons as measured by the High Energy Particle Spectrometer on the Upper Atmosphere Research Satellite and changes in the chemistry of the mesosphere as measured by CLAES and MLS on the same spacecraft. The following tasks are excerpted from the statement of work:

1. Using data from the PEM/HEPS instrument aboard the UARS spacecraft, periods of highly relativistic electron precipitation will be identified, and Spaceraid will join in the selection of specific periods for more detailed analysis.
2. Spaceraid will provide energy spectra and spatial and temporal distributions of precipitating relativistic electron fluxes based on the PEM/HEPS data for the specific event periods selected in task #1. The results will be provided to distinguish stably trapped, drift loss cone, and bounce loss cone fluxes.
3. Spaceraid will contribute to the analysis, presentation, and publication of the initial results of this project.

Task 1

From continuous survey plots of HEPS data, the daily peak flux of trapped electrons with energies > 2 MeV were surveyed for the entire database up to March 1995 and three time periods showing large changes in flux within two days or less were selected for further study. The periods chosen within the operational lifetime of CLAES as an initial criterion were: 1. 8 - 9 November 1991 when a large magnetic storm precipitated the bulk of the relativistic electrons seen at low altitude
2. 16 - 17 December 1991, a modest relativistic electron event
3. 10 - 20 May 1992, the largest relativistic electron flux event in the database

Claes viewed southward from the spacecraft during the first period and northward during the second and third periods. The magnetic local time during the times of intense electron precipitation was between ~0900 and ~1600 hours for all three periods.

Task 2


Spatial and temporal distributions of the energy spectra of precipitating relativistic electrons were extracted from the HEPS data for the time periods selected in Task 1. From these, ion production versus altitude profiles were calculated using a code furnished by C. H. Jackman (NASA, Goddard Space Flight Center). Examples of the spectra and corresponding ion production profiles are attached.

Task 3


For the highest intensity relativistic electron flux event of May 1992, global plots of trapped as well as precipitating electrons $E > 2$ MeV were prepared. Detailed spectra were extracted and ion production profiles were calculated during the period for the southern hemisphere region for correlation with HALOE measurements as well as for the drift loss cone region in the north to compare with CLAES and MLS atmospheric measurements.

Since the precipitating relativistic electron fluxes in the initially selected data periods were in sunlit conditions making the atmospheric chemistry changes due to the electrons perhaps less discernible from those produced by solar UV, the events with the most pronounced flux changes are the most promising. The rapid removal of relativistic electron fluxes on Nov. 8, 1991 would allow observation of ozone replenishment while the buildup of the fluxes between May 10 and 12, 1992 presents the largest change in ionization seen at altitudes below 60 km in the north and resulting in ozone depletion. The ion profiles due to precipitating electrons show five orders of magnitude increase at nearly the same location between May 10 and May 12, 1992 with the intensity of the ion production at 60 km equal to 10^2 ion-pairs/cm³-sec.


PRECIPITATING ELECTRON SPECTRA, 8 - 9 NOVEMBER 1991, ANOMALY/SOUTH




ION PRODUCTION PROFILES, 8 - 9 NOV., 16 - 17 DEC. 1991, 8 & 12 MAY 1992




PRECIPITATING ELECTRON SPECTRA, MAY 1992 - DRIFT LOSS CONE / NORTH



ION PRODUCTION PROFILES, MAY 1992 - DRIFT LOSS CONE (180 - 300 E / NORTH)



HEPS PRECIPITATING ELECTRONS, MAY 1992 - ANOMALY, SOUTH



ION PRODUCTION PROFILES, 10-18 MAY 1992 - ANOMALY / SOUTH

